

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : <b>A61K 31/59</b>	<b>A1</b>	(11) International Publication Number: <b>WO 94/16711</b> (43) International Publication Date: <b>4 August 1994 (04.08.94)</b>
<p>(21) International Application Number: <b>PCT/US94/01172</b></p> <p>(22) International Filing Date: <b>31 January 1994 (31.01.94)</b></p> <p>(30) Priority Data: <b>08/010,823</b>      <b>29 January 1993 (29.01.93)</b>      <b>US</b></p> <p>(71) Applicant: <b>LUNAR CORPORATION [US/US]; 313 West Belkline Highway, Madison, WI 53713 (US).</b></p> <p>(72) Inventors: <b>GULBRANDSEN, Carl, E.; 1506 Wood Lane, Madison, WI 53705 (US). MOSS, Richard, L.; 1326 Boundary Road, Middleton, WI 53562 (US).</b></p> <p>(74) Agents: <b>GULBRANDSEN, Carl, E. et al.; Stroud, Stroud, Willink, Thompson &amp; Howard, Suite 300, 25 West Main Street, P.O. Box 2236, Madison, WI 53701-2236 (US).</b></p>		<p>(81) Designated States: <b>CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</b></p> <p><b>Published</b> <i>With international search report.</i></p>
<p>(54) Title: <b>PREVENTION AND TREATMENT OF MYOCARDIAL FAILURE</b></p> <p>(57) Abstract</p> <p>Method of increasing the strength of contraction in the mammalian heart muscle by administering to the mammal an effective amount of an activated vitamin D compound, i.e., a 1<math>\alpha</math>-hydroxylated vitamin D compound which binds with the vitamin D receptor and produces a positive inotropic effect in the heart muscle. The activated vitamin D compound may be given as a means to prevent myocardial failure or to treat myocardial failure.</p>		

BEST AVAILABLE COPY

M3

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Martinique
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Cote d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**PREVENTION AND TREATMENT OF MYOCARDIAL FAILURE****TECHNICAL FIELD**

5           This invention relates a method of treating myocardial failure, more specifically it relates to the use of active forms of vitamin D to increase the strength contraction of the heart muscle.

**BACKGROUND OF THE INVENTION**

10           Heart failure is a common clinical condition and results in a significant morbidity and mortality. It is defined as the pathophysiologic state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate  
15           commensurate with the requirements of the metabolizing tissues or can do so only from an abnormally elevated filling pressure. Heart failure is frequently, but not always, caused by a defect in myocardial contraction wherein the strength of contraction of the heart muscle  
20           is diminished. In such a case, the term myocardial failure is appropriate. Few therapies exist for myocardial failure that are effective and do not present significant undesirable side effects. The most common treatment of myocardial failure is the administration of  
25           cardiac glycosides such as digitalis. While digitalis can alleviate the symptoms and improve cardiac hemodynamics in heart failure, it, as well as the other cardiac glycosides, has a low margin of safety. Such potent drugs cause cardiac dysrhythmias and neurological  
30           problems as well as nausea, abdominal pain and headache. Further, drug interaction problems are reported with the cardiac glycosides and other common drugs.

35           What is needed is a method of increasing the strength of the heart contraction without the above-described undesirable side effects.

-2-

## DETAILED DESCRIPTION

The present invention is for a method of treating myocardial failure using an active form of vitamin D. Vitamin D is known to be important in the regulation of calcium metabolism in animals and man. See, Harrison's  
5 Principals of Internal Medicine: Part Eleven, "Disorders of Bone and Mineral Metabolism", Chapter 335, E. Braunwald et al., (eds.), McGraw-Hill, New York, 1987, pp. 1860-1865.

10 It is known that vitamin D<sub>3</sub> must be hydroxylated in the 1 and the 25 position before it is activated, i.e., before it will produce a biological response. A similar metabolism appears to be required to activate the other forms of vitamin D, e.g., vitamin D<sub>2</sub> and vitamin D<sub>4</sub>. As  
15 is generally understood and used herein, the term "vitamin D" is intended to include vitamins D<sub>3</sub>, D<sub>2</sub>, and D<sub>4</sub>. The term activated vitamin D, as used herein, is intended to refer to vitamin D which has been hydroxylated in at least the 1 position of the A ring  
20 and binds with the vitamin D receptor, e.g., 1,25-dihydroxyvitamin D<sub>3</sub>.

The 1 $\alpha$ -hydroxyvitamin D of the present invention has the general formula described in Figure I wherein A  
25 and B are either hydrogen or a carbon-to-carbon bond, thus forming a double bond between C22 and C23, R<sub>2</sub>, and R<sub>3</sub> can be either hydrogen, hydroxy, lower alkyl, O-lower alkyl, O-lower acyl, O-aromatic acyl or fluoro, and where R<sub>4</sub> is hydrogen or lower alkyl along with an acceptable excipient.

30 In the formulae shown in this specification and in the claims a wavy line to substituent X indicates that the substituent can be either  $\alpha$  or  $\beta$  stereoisomeric form. Wherever in this specification and in the claims the word "lower" is used as a modifier of alkyl or acyl  
35 it is intended to identify a hydrocarbon chain having from about 1 to 4 carbon atoms and can be either a straight chain or branched chain configuration. Specific examples of such hydrocarbon chains are:

-3-

methy1, ethyl, propyl, butyl, isobutyl or t-butyl, and  
 formyl, acetyl, propionyl, or butyryl. The word  
 "aromatic acyl" as used herein and in the claims is  
 meant to identify a benzoyl group or a substituted  
 5 benzoyl group such as nitrobenzoyl or dinitrobenzoyl.




Figure I

Among the preferred active vitamin D compounds are:

- 1 $\alpha$ ,25-dihydroxy-cholecalciferol[1 $\alpha$ ,25-(OH) $_2$ D $_3$ ]
- 1 $\alpha$ -hydroxy-cholecalciferol[1 $\alpha$ -(OH)D $_3$ ]
- 1 $\alpha$ ,24-dihydroxy-cholecalciferol[1 $\alpha$ ,24-(OH) $_2$ D $_3$ ]
- 10 1 $\alpha$ ,25-dihydroxy-ergocalciferol[1 $\alpha$ ,25-(OH) $_2$ D $_2$ ]
- X 1 $\alpha$ -hydroxy-ergocalciferol[1 $\alpha$ -(OH)D $_2$ ]
- X 1 $\alpha$ ,24(s)-dihydroxy-ergocalciferol[1 $\alpha$ ,24(s)-(OH) $_2$ D $_2$ ]
- 1 $\alpha$ ,25-dihydroxy-vitamin D $_4$ [1 $\alpha$ ,25-(OH) $_2$ D $_4$ ]
- X 1 $\alpha$ -hydroxy-vitamin D $_4$ [1 $\alpha$ -(OH)D $_4$ ]
- 15 X 1 $\alpha$ ,24-dihydroxy-vitamin D $_4$ [1 $\alpha$ ,24-(OH) $_2$ D $_4$ ]

The above-described active forms of vitamin D can  
 be prepared as described in U.S. Patents 3,993,675;  
 4,022,891; 4,195,027; 4,234,495; 4,508,651 and co-  
 pending U.S. applications 07/940,246 and 07/991,493 all  
 20 incorporated herein by reference.

Functionally, vitamin D is more appropriately  
 considered a hormone than a vitamin. When activated,  
 vitamin D interacts with a vitamin D receptor protein  
 and this interaction ultimately results in some form of

-4-

biological response. For example,  $1\alpha,25$ -dihydroxyvitamin  $D_3$  is known to be a potent stimulator of calcium absorption from the intestine which is mediated by the interaction of the  $1\alpha,25$ -dihydroxyvitamin  $D_3$  molecule and the vitamin D receptor protein located in the epithelial cells (enterocytes) which line the intestine.

In recent years it has become evident that the vitamin D receptor protein is widely distributed in the bodies of animals and man. Thus, it is not surprising that besides calcium homeostasis, activated vitamin D has been implicated in osteogenesis, modulation of immune response, modulation of the process of insulin secretion by the pancreatic B cell, muscle cell function and the differentiation and growth of epidermal and hemopoietic tissues.

More recently,  $1\alpha,25$ -dihydroxyvitamin  $D_3$  receptors have been shown to exist in the rat heart, (Walters et al., J. Mol. Cell Cardiol. 18: 67-72 (1986)) and this has prompted the speculation that vitamin D may play a role in cardiac function. Until the present invention, the prevailing view, which was based on studies of cardiac hemodynamics in vitamin  $D_3$  deficient rats, was that  $1\alpha,25$ -dihydroxyvitamin  $D_3$  produced a direct negative inotropic effect in the heart, presumably by promoting the sequestering of calcium in the myocardium. (Weishaar and Simpson, Am. J. Physiol. 253 (Endocrinol. Metab. 16): E675-E683 (1987)).

Contrary to the hypothesis of Weishaar and Simpson, the present inventors have found that active forms of vitamin  $D_3$ , including  $1\alpha,25$ -dihydroxyvitamin  $D_3$ , produce a direct positive inotropic effect in the mammalian myocardium, i.e., increases the strength of the contraction of the heart muscle.

#### Example 1: Positive Inotropic Effect

Rat right ventricular papillary muscles were mounted in an experimental chamber and stimulated at

-5-

0.3-0.7 Hz with a single pulse, broad field stimulation via platinum plate electrodes. The preparation was continuously perfused at 22-24° with oxygenated modified Tyrode's solution, pH 7, containing 2 mM  $\text{Ca}^{2+}$ . Twitch tension of the preparation was measured by suturing one end of the preparation to a force transducer and the other end to a three-way positioner. Muscles attaining a stable baseline twitch tension were then perfused with 0.1 to 6.25  $\mu\text{M}$  of  $1\alpha,25$ -dihydroxyvitamin  $\text{D}_3$ . In nine experiments  $1\alpha,25$ -dihydroxyvitamin  $\text{D}_3$  increased steady-state twitch tension an average  $14 \pm 11\%$  (range of 4-41%). The effects of  $1\alpha,25$ -dihydroxyvitamin  $\text{D}_3$  were reversed by drug washout. These results indicate that  $1\alpha,25$ -dihydroxyvitamin  $\text{D}_3$  has a positive inotropic effect on the mammalian myocardium.

**Example 2: Prevention of Congestive Heart Failure**

An oral dosage formulation containing  $1\alpha,25$ -dihydroxyvitamin  $\text{D}_3$  is evaluated in a double blind study for efficacy in preventing the development of heart failure caused by myocardial failure. The formulation evaluated contains 0.25  $\mu\text{g}$  of  $1\alpha,25$ -dihydroxyvitamin  $\text{D}_3$ . The control formulation is identical except that it does not contain the  $1\alpha,25$ -dihydroxyvitamin  $\text{D}_3$ . Five hundred normal subjects between the ages of 55 and 65 are selected. The subjects are divided into an experimental and control population. They are instructed to take the medication twice a day, in the morning and in the evening.

Evaluations of cardiovascular hemodynamics are conducted at six-month intervals by a physician. The final evaluation is carried out at the end of three years of preventive therapy. The results of the study show that daily oral administration of  $1,25$ -dihydroxyvitamin  $\text{D}_3$  significantly reduces the occurrence of myocardial failure in the experimental group as compared with the control.

-6-

As the above example illustrates, preventive benefit in reducing the occurrence of myocardial failure is derived from daily administration of a relatively low dosage of  $1\alpha,25$ -dihydroxyvitamin  $D_3$ . For treatment purposes, however, a higher dosage would be desired. However, the vitamin  $D_3$  compounds, particularly,  $1\alpha,25$ -dihydroxyvitamin  $D_3$  cannot safely be administered at a dosage greater than  $1.0 \mu\text{g}$  per day without causing hypercalcemia and hypercalciuria in a large portion of the population. In that regard, the active forms of vitamin  $D_2$  and vitamin  $D_4$  are more suitable for, while they display a high binding activity with respect to the vitamin D receptor, they have a much lower calcemic effect and are thus much less toxic. See, for example, co-pending U.S. application 07/940,246 which is incorporated herein by reference. Preferred in this regard are  $1\alpha$ -hydroxy-ergocalciferol [ $1\alpha$ -(OH) $D_2$ ],  $1\alpha,24$ (s)-dihydroxy-ergocalciferol [ $1\alpha,24$ (s)-(OH) $_2D_2$ ],  $1\alpha$ -hydroxy-vitamin  $D_4$  [ $1\alpha$ -(OH) $D_4$ ] and  $1\alpha,24$ -dihydroxy-vitamin  $D_4$  [ $1\alpha,24$ -(OH) $_2D_4$ ].

Advantageously, the vitamin  $D_2$  and  $D_4$  compounds of the present invention or combinations thereof with other therapeutic agents can be administered in dosage amounts of from 0.1 to 10.0 micrograms per day. These compounds can be administered as sterile parenteral solutions by injection or intravenously or by alimentary canal in the form of oral dosages, or by suppository. In relation to treatment of early stage myocardial failure doses from about 1.5 to about 6.0 micrograms per day are generally effective. For more advanced stages of myocardial failure, it may be advisable to administer the compounds of the present invention in conjunction with more traditional therapies such as the cardiac glycosides. Surprisingly, it is found that the compounds of the present invention produce a synergistic response when administered in conjunction with another positive inotropic compound such as the glycosides. This synergistic effect allows the physician to administer a



-7-

lower dosage of the glycosides and helps to avoid many of the undesirable side effects of the glycosides. If the compounds of the present invention are administered in combination with other therapeutic agents, the proportions of each of the compounds in the combination being administered will be dependent on the particular agents being used and the degree of heart failure being treated. It being understood that the specific dosage administered in any given case will be adjusted in accordance with the specific compounds being administered, the stage of the myocardial failure to be treated, the condition of the subject and the other relevant medical facts that may modify the activity of the drug or the response of the subject, as is well known by those skilled in the art.

While the present invention has now been described and exemplified with some specificity, those skilled in the art will appreciate the various modifications, including variations, additions, and omissions, that may be made in what has been described. Accordingly, it is intended that these modifications also be encompassed by the present invention and that the scope of the present invention be limited solely by the broadest interpretation that lawfully can be accorded the appended claims.

-8-

## CLAIMS:

1. A method for preventing or treating myocardial failure in a mammal comprising administering to said mammal an effective amount of a compound of the general structure of Figure I wherein A and B are either hydrogen or a carbon-to-carbon bond thus forming a double bond between C22 and C23, R<sub>1</sub> and R<sub>2</sub> can be either hydrogen, hydroxy, lower alkyl, O-lower alkyl, O-lower acyl, O-aromatic acyl or fluoro, and where R<sub>4</sub> is hydrogen or lower alkyl along with an acceptable excipient.




Figure I

2. A method for preventing or treating myocardial failure as described in claim 1 wherein the compound of Figure I is selected from the group consisting of 1 $\alpha$ ,25-dihydroxycholecalciferol, 1 $\alpha$ -hydroxycholecalciferol, 1 $\alpha$ ,24-dihydroxycholecalciferol, 1 $\alpha$ ,25-dihydroxy-ergocalciferol, 1 $\alpha$ -hydroxy-ergocalciferol, 1 $\alpha$ ,24(s)-dihydroxy-ergocalciferol, 1 $\alpha$ ,25-dihydroxy-vitamin D<sub>3</sub>, 1 $\alpha$ -hydroxy-vitamin D<sub>3</sub>, 1 $\alpha$ ,24-dihydroxy-vitamin D<sub>3</sub>.

3. A method for increasing the strength of the contraction in the heart of a mammal comprising

-9-

administering to said mammal an effective amount of a compound of the general structure of Figure I wherein A and B are either hydrogen or a carbon-to-carbon bond thus forming a double bond between C22 and C23, R<sub>1</sub>, and R<sub>2</sub> can be either hydrogen, hydroxy, lower alkyl, O-lower alkyl, O-lower acyl, O-aromatic acyl or fluoro, and where R<sub>3</sub> is hydrogen or lower alkyl along with an acceptable excipient.




Figure I

4. A method for increasing the strength of contraction in the heart of a mammal as claimed in claim 3 wherein the compound of Figure I is selected from the group consisting of 1 $\alpha$ ,25-dihydroxycholecalciferol, 1 $\alpha$ -hydroxy-cholecalciferol, 1 $\alpha$ ,24-dihydroxy-cholecalciferol, 1 $\alpha$ ,25-dihydroxy-ergocalciferol, 1 $\alpha$ -hydroxy-ergocalciferol, 1 $\alpha$ ,24(s)-dihydroxy-ergocalciferol, 1 $\alpha$ ,25-dihydroxy-vitamin D<sub>3</sub>, 1 $\alpha$ -hydroxy-vitamin D<sub>3</sub>, 1 $\alpha$ ,24-dihydroxy-vitamin D<sub>3</sub>.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/01172

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(S) :A61K 31/59

US CL :514/167, 168

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/167, 168

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, REGISTRY (STRUCTURE), CA, MEDLINE

search terms: heart, myocardial, infarction, disease, disorder, damage

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 4,501,737 (YAMOTO ET AL) 26 FEBRUARY 1985, see entire document.	1-4
A	Chemical Abstracts, Volume 112, No. 15, issued 1990, Weishaar et al, "Involvement of vitamin D3 with cardiovascular function. Effects on physical and morphological properties" see page 148, column 2, abstract 133090g, Am. J. Physiol., 258(1, Pt. 1), E134-E142.	1-4

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be part of particular relevance	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	A*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*T* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 12 APRIL 1994	Date of mailing of the international search report 04 MAY 1994
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer WILLIAM JARVIS
Facsimile No. (703) 305-3230	Telephone No. (703) 308-1235

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**